



A Reliable Research Partner in Life Science and Medicine

## Recombinant Human BMP-5 protein(His Tag)

Catalog Number: PKSH034131

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$	escrip				
	00	OP	m		٦n
J		VI.		w	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

Species Human

Source E.coli-derived Human BMP-5 protein Ala 317-His 454, with an C-terminal His

Calculated MW16.6 kDaObserved MW17 kDaAccessionP22003

**Bio-activity** Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The

ED<sub>50</sub> for this effect is  $< 0.17 \mu g/mL$ .

**Properties** 

**Purity** > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Background

Growth factor of the TGF-beta superfamily that plays essential roles in many developmental processes, including cartilage and bone formation or neurogenesis. Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR. In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes.

Can also signal through non-canonical pathway such as MAPK p38 signaling cascade to promote chondrogenic differentiation.