A Reliable Research Partner in Life Science and Medicine

# Recombinant Human IL-20RA/IL-20R1 Protein (Fc Tag)

Catalog Number: PKSH032575

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$		crip					
	00	0	РΤ	n	ŤΤ		m
v	$\mathbf{c}$	v.		w	w	w	ш

**Species** Human

Source HEK293 Cells-derived Human IL-20RA; IL-20R1 protein Val30-Lys250, with an C-

terminal Fc

Calculated MW52.4 kDaObserved MW70-85 kDaAccessionQ9UHF4

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Interleukin-20 Receptor Subunit  $\alpha$  (IL20RA) is a single-pass type I membrane protein that is a member of the type II cytokine receptor family. IL20RA is synthetized a 553 amino acid glycoprotein precursor containing a 29 amino acid signal peptide; a 221 amino acid extracellular domain with two fibronectin type-III domains; a 24 amino acid transmembrane region; and a 279 amino acid intracellular domain. IL20RA is widely expressed with highest levels found in skin and testis and high levels in brain. IL20RA forms a heterodimer with IL20RB; and the complex serves as a receptor for IL19; IL20 and IL24. IL20RA also forms a heterodimer with the unique and specific receptor IL10RB and functions as the receptor for IL26.

### For Research Use Only