

## APC Anti-Mouse CD226 Antibody[10E5]

Catalog Number: E-AB-F1402E

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2b, κ
<b>Clone No.</b>	10E5
<b>Isotype Control</b>	APC Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842E]
<b>Conjugation</b>	APC
<b>Conjugation Information</b>	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	DNAM-1;PTA1(platelet and T cell activation antigen 1);TLISA1;LFA-1 associated Molecule PTA-1
<b>Uniprot ID</b>	Q8K4F0
<b>Gene ID</b>	225825
<b>Background</b>	CD226 (DNAM-1) is constitutively expressed on native CD8+ cells and on CD4+ T cells, macrophages and NK cells.This antibody (10E5) was reported to bind about 40% of inactivated CD4+ cells and binds only to differentiated Th1 cells, but not to Th2 or Th0 cells.It is also reported to suppress antigen-specific T cell expansion and EAE ( experimental allergic encephalitis) mediated by Th1 cells.

### For Research Use Only