# **Elabscience Biotechnology Co., Ltd.**



A Reliable Research Partner in Life Science and Medicine

# **VWA5A Polyclonal Antibody**

catalog number: E-AB-10998

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Reactivity** Human

Immunogen Recombinant protein of human VWA5A

Host Rabbit Isotype IgG


Purification Affinity purification
Conjugation Unconjugated


**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

**IHC** 1:100-1:300

#### Data





Immunohistochemistry of paraffin-embedded Human cervical Immunohistochemistry of paraffin-embedded Human lung cancer tissue using VWA5A Polyclonal Antibody at dilution cancer tissue using VWA5A Polyclonal Antibody at dilution 1:80

#### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

VWA5A (von Willebrand factor A domain containing 5A), also known as BCSC-1 (breast cancer suppressor candidate 1) or LOH11CR2A (loss of heterozygosity 11 chromosomal region 2 gene A protein), is a 786 amino acid protein containing one VIT domain and one VWFA domain. VWA5A is expressed at low levels in various tissues, with no expression found in 80% of tumor cell lines. Likely acting as a tumor suppressor gene, deletion of VWA5A leads to loss of heterozygosity (LOH) in breast and ovarian tumors, and may have an important role as a potential gene therapy targe t. Abnormal expression of VWA5A may lead to an increase in adhesion of CNE-2L2 cells associated with an increase in expression of E-cadherin, alpha-catenin, and p53, resulting in a decrease of malignant activity in cells with ectopic expression of VWA5A.

## For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn