


Recombinant SUCLG1 Monoclonal Antibody

catalog number: AN302017L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	Rabbit IgG
Clone	C321638
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:500-1:10000
IHC	1:500-1:1000

Data





Western Blot with SUCLG1 Monoclonal Antibody at dilution of 1:10000. Lane 1: HepG2

Observed-MW:36-26 kDa
Calculated-MW:36 kDa




Western Blot with SUCLG1 Monoclonal Antibody at dilution of 1:1000. Lane 1: A431

Observed-MW:36-26 kDa
Calculated-MW:36 kDa



Western Blot with SUCLG1 Monoclonal Antibody at dilution of 1:2000. Lane 1: HeLa

Observed-MW:36-26 kDa
Calculated-MW:36 kDa



Immunohistochemistry of paraffin-embedded Human kidney using SUCLG1 Monoclonal Antibody at dilution of 1:1000.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Immunohistochemistry of paraffin-embedded Human
liver cancer using SUCLG1 Monoclonal Antibody at
dilution of 1:1000.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Succinyl-CoA synthetase α subunit (SUCLG1) catalyzes the conversion of succinate to succinyl-CoA and plays a key role in the citric acid cycle. Deficiency of this enzyme leads to a variety of diseases including fatal infantile lactic acidosis and mitochondrial hepatoencephalomyopathy.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2