



A Reliable Research Partner in Life Science and Medicine

Human S100A12 Antibody Pair Set

Catalog No. E-KAB-0529 Applications ELISA

Synonyms CAAF1;CAGC;CGRP;ENRAGE;MRP-6;MRP6;p6;Calgranulin C

Kit components & Storage

Title	Specifications	Storage
Human S100A12 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Human S100A12 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0529)		
		Human S100A12 Capture Antibody	Human S100A12 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human S100A12	Recombinant Human S100A12 protien	
Information		protien		
	Swissprot	P80511		
Product details	Reactivity	Human	Human	
	Host	Rabbit	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Antigen Affinity	Antigen Affinity	
Specificity		Detects Human S100A12 in ELISAs.		

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn





A Reliable Research Partner in Life Science and Medicine

Applications

Human S100A12 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human S100A12 Capture	
Capture		Antibody	10
			Aige
ELISA	1:1000-1:10000	Human S100A12	Optical Density
Detection		Detection Antibody	O 0.1
		(Biotin)	
			0.01

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calciumbinding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein is proposed to be involved in specific calcium-dependent signal transduction pathways and its regulatory effect on cytoskeletal components may modulate various neutrophil activities.

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn