Cleaved-CASP4 p20 (Q81) Polyclonal Antibody

Catalog Number: E-AB-30021



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		- 40	
Desc	\mathbf{r}	$\mathbf{n}\mathbf{T}\mathbf{I}$	\mathbf{on}
DUSU	ш		ULL

Reactivity Human

Immunogen Synthesized peptide derived from the Internal region of human Caspase-4 p20

Host Rabbit
Isotype IgG

Purification Affinity purification

Conjugation Unconjugated

Formulation PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4

A	oplications	Recommended Dilution

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000

Data



Western Blot analysis of 3T3 cells using Cleaved-CASP4 p20 (Q81) Polyclonal Antibody at dilution of 1:2000.

Observed MW:43+22kDa Calculated Mw:43kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a member of the cysteine proteases that plays important roles in apoptosis, cell migration and the inflammatory response. The encoded protein mediates production of pro-inflammatory cytokines by macrophages upon bacterial infection. Mice lacking the encoded protein are resistant to endotoxic shock induced by lipopolysaccharide. A 5-bp deletion encompassing a splice acceptor junction resulting in alternate splicing and a shorter non-functional isoform in certain mouse strains has been described. Although its official nomenclature is "caspase 4, apoptosis-related cysteine peptidase", this gene and its encoded protein have historically been called caspase 11. This gene is present in a cluster of three caspase genes on chromosome 9.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com