

Recombinant Human CLIC4 Protein (His Tag)

Catalog Number:PKSH030859



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CLIC4L;H1;huH1;MTCLIC;p64H1
Species	Human
Expression Host	E.coli
Sequence	Ala 2-Lys 253
Accession	Q9Y696-1
Calculated Molecular Weight	29.6 kDa
Observed molecular weight	30 kDa
Tag	N-His

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Chloride intracellular channel protein 4; also known as Intracellular chloride ion channel protein p64H1 and CLIC4; is a member of the chloride channel CLIC family. It contains oneGST C-terminal domain. CLIC4 is a member of a family of intracellular chloride channels. It is regulated by p53; c-Myc; and tumor necrosis factor-alpha. CLIC4 is detected in epithelial cells from colon; esophagus and kidney (at protein level). CLIC4 has alternate cellular functions like a potential role in angiogenesis or in maintaining apical-basolateral membrane polarity during mitosis and cytokinesis. CLIC4 could promote endothelial cell proliferation and regulate endothelial morphogenesis (tubulogenesis). Expression of CLIC4 is prominent in heart; kidney; placenta and skeletal muscle. Overexpression of CLIC4 in cancer cells inhibits tumor growth. Conversely; overexpression of CLIC4 in tumor stromal cells stimulates tumor growth. Thus; CLIC4 participates in normal

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CLIC4 Protein (His Tag)

Catalog Number:PKSH030859



and pathological processes and may serve as a useful target for therapies in disturbances of homeostasis and neoplastic transformation. Loss of CLIC4 in tumor cells and gain in tumor stroma is common to many human cancers and marks malignant progression. Up-regulation of CLIC4 in tumor stroma is coincident with myofibroblast conversion; generally a poor prognostic indicator. Reactivation and restoration of CLIC4 in tumor cells or the converse in tumor stromal cells could provide a novel approach to inhibit tumor growth.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017