# **Elabscience**®

## **MAGED1** Polyclonal Antibody

### catalog number: E-AB-10444

Note: Centrifuge before opening to ensure complete recovery of vial contents.


1:25-1:100

| Description  |  |
|--------------|--|
| Reactivity   | Human;Mouse;Rat  |
| Immunogen    | Recombinant protein of human MAGED1  |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Affinity purification  |
| Buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution   |
| WB           | 1:500-1:2000   |

### Data


WB

IHC



Western Blot analysis of A549, Jurkat and 293T cell using MAGED1 Polyclonal Antibody at dilution of 1:1200

#### Calculated-MW:86 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer using MAGED1 Polyclonal Antibody at dilution of

1:30

Immunohistochemistry of paraffin-embedded Human colon cancer using MAGED1 Polyclonal Antibody at dilution of 1:30

**Preparation & Storage** Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage The product is shipped with ice pack, upon receipt, store it immediately at the Shipping temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

# **Elabscience**®

This gene is a member of the melanoma antigen gene (MAGE) family. Most of the genes of this family encode tumor specific antigens that are not expressed in normal adult tissues except testis. Although the protein encoded by this gene shares strong homology with members of the MAGE family, it is expressed in almost all normal adult tissues. This gene has been demonstrated to be involved in the p75 neurotrophin receptor mediated programmed cell death pathway. Three transcript variants encoding two different isoforms have been found for this gene.

For Research Use Only