

Recombinant Human KIAA0101/p15/PAF Protein (His Tag)

Catalog Number: PKSH031315

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | E.coli-derived Human KIAA0101/p15/PAF protein Met 1-Glu 111, with an N-terminal His |
| Calculated MW | 13.8 kDa |
| Observed MW | 19 kDa |
| Accession | NP_055551.1 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 97 % as determined by reducing SDS-PAGE. |
| Endotoxin | Please contact us for more information. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

KIAA0101, also known as p15(PAF), is a proliferating cell nuclear antigen-associated factor which interacts with proliferating cell nuclear antigen(PCNA). It was initially isolated in a yeast two-hybrid screen for PCNA binding partners, and was shown to bind PCNA competitively with the cell cycle regulator p21(WAF). KIAA0101 is localized primarily in the nucleus. It shares the conserved PCNA binding motif with several other PCNA binding proteins including CDK inhibitor p21. KIAA0101 is involved in cell proliferation and plays a role in early tumor recurrence (ETR), and prognosis of hepatocellular carcinoma (HCC). KIAA0101 is expressed predominantly in liver, pancreas and placenta. It cannot be detected in heart or brain. It is highly expressed in a number of tumors, especially esophageal tumors, in anaplastic thyroid carcinomas and in non-small-cell lung cancer lines. Overexpression of KIAA0101 predicts high stage, early tumor recurrence, and poor prognosis of hepatocellular carcinoma. It also may be involved in protection of cells from UV-induced cell death.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017