

0.25% Trypsin Solution, with EDTA, phenol red, dissolved in PBS

Cat. No. : PB180229

Size: 100mL

General Information

Product Form	Liquid
Concentration	0.0025
HEPES	Negative
Phenol red	10 mg/L
Trypsin	2500 mg/L
EDTA • 2Na	200 mg/L
Storage	-5~-20°C, Shading Light
Shipping	Ice bag
Expiration date	12 months

Background

Trypsin is a serine hydrolase that can cut the base-side segment of the lysine and arginine residue in the polypeptide chain, hydrolyze the protein between the cells, and destroy the connection between the cells, so that the tissue or the adherent cells are dispersed into a single cell. The activity of the trypsin is related to the characteristics of the tissue or the cells, the concentration of the trypsin, the temperature and the time of action. At pH 8.0 and 37°C, the effect of the trypsin is the best. Therefore, the concentration of trypsin, the temperature and the time of the pancreatic enzyme should be grasped to avoid the excessive damage of the cells.

In general, the frequently-used working concentration of the trypsin is 0.25%, while the concentration of trypsin for the semi-adherent cells or the cells is 0.05%. Since EDTA is able to bind Ca^{2+} and Mg^{2+} to destroy the cell connection to promote the dissociation of the cells, a certain amount of EDTA is often added to the trypsin solution to enhance the dissociation effect

Notes

1. This product is for research use only.
2. This product is sterilized by 0.1μm filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the pollution.
4. It is not recommended to store this product at room temperature for a long time.