

## Recombinant Mouse IGFBP-2/IGFBP2 Protein (Fc Tag)

**Catalog Number:** PKSM040345

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse IGFBP-2/IGFBP2 protein Met1-Gln305, with an C-terminal hFc
<b>Calculated MW</b>	56.6 kDa
<b>Accession</b>	NP_032368.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

IGFBP-2, also known as IGFBP2, is a insulin-like growth factor-binding protein (IGFBP). IGFBPs prolong the half-life of the IGFs , control bioavailability, activity, and distribution of insulin-like growth factor (IGF) through high-affinity IGFBP/IGF complexes. Six high-affinity IGF-binding proteins (IGFBP-1 to -6) have been identified. The six IGFBPs are structurally related but encoded by distinct genes. IGFBPs have a high affinity for IGFs. Some members of the IGFBP family have been consistently shown to inhibit IGF actions by preventing them from gaining access to the IGF receptors, while others potentiate IGF actions by facilitating the ligand-receptor interaction. IGFBP-2 is overexpressed in many malignancies and is often correlated with an increasingly malignant status of the tumor, pointing to a potential involvement of IGFBP-2 in tumorigenesis. It contains 1 IGFBP N-terminal domain and 1 thyroglobulin type-1 domain. It inhibits IGF-mediated growth and developmental rates.