

Recombinant Human Serpin A7/TBG (C-6His)

Catalog Number: PKSH033879

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Serpin A7;TBG protein Ala21-Ala415, with an C-terminal His
Calculated MW	49.9 kDa
Observed MW	55-65 kDa
Accession	P05543
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Serpin A7 is a secreted protein which belongs to the Serpin family. As a 54 kDa protein, Serpin A7 is synthesized primarily in the liver and secreted in plasma. Genomically, Serpin A7 is a serpin, although it has no inhibitory function like many other members of this class of proteins. Serpin A7 binds thyroid hormone in circulation. It is one of three proteins (along with transthyretin and albumin) responsible for carrying the thyroid hormones thyroxine (T4) and 3,5,3'-triiodothyronine (T3) in the bloodstream. Of these three proteins, Serpin A7 has the highest affinity for T4 and T3, but is present in the lowest concentration. Despite its low concentration, Serpin A7 carries the majority of T4 in serum. Due to the very low serum concentration of T4 and T3 in the blood, Serpin A7 is rarely more than 25% saturated with its ligand. Unlike transthyretin and albumin, Serpin A7 has a single binding site for T4/T3. Serpin A7 tests are sometimes used in finding the reason for elevated or diminished levels of thyroid hormone, and defects in Serpin A7 are a cause of Serpin A7 deficiency.