

Recombinant Human ADP-ribosyl Cyclase/cyclic ADP-ribose Hydrolase 1/CD38 (N-6His-Avi) Biotinylated

Catalog Number: PKSH033880

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Human |
| Source | HEK293 Cells-derived Human CD38 protein Val43-Ile300, with an N-terminal His & Avi |
| Calculated MW | 33.3 kDa |
| Observed MW | 40-50 kDa |
| Accession | P28907 |
| Bio-activity | Not validated for activity |

Properties

| | |
|----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Concentration | Subject to label value. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles. |
| Shipping | This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C. |
| Formulation | Supplied as a 0.2 µm filtered solution of PBS, pH7.4. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD38, also known as ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, is a Signal-anchor for type II membrane protein. CD38 is able to transform NAD⁺ to ADP-D-ribose and nicotinamide. It also can transform NADP⁺ to nicotinate-adenine dinucleotide phosphate and nicotinamide. CD38 is expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma. Synthesizes the second messengers cyclic ADP-ribose and nicotinate-adenine dinucleotide phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017