Recombinant Mouse TRAIL R2/TNFRSF10B Protein (AVI & His Tag)

Catalog Number: PKSM041369



Note: Centrifuge before opening to ensure complete recovery of vial contents.

			4.0
1116	100	rin	tion
\mathbf{v}		TID	tion

 Species
 Mouse

 Mol_Mass
 16.4 kDa

 Accession
 O9OZM4

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Mouse tumor necrosis factor receptor superfamily member 10B (TNFRSF10B) is a member of the TNFR family which contains 1 death domain and 3 TNFR-Cys repeats. TNFRSF10B exhibits high structural and functional homology to TRAIL-R1 (DR-4). TNFRSF10B is highly expressed in heart, lung, lymphocytes, spleen and kidney. In addition, it is regulated by the tumor suppressor p53. TNFRSF10B is the receptor for the cytotoxic ligand TNFSF10/TRAIL. It promotes the activation of NF-kappa-B and is essential for ER stress-induced apoptosis. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis.

For Research Use Only