



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Human CD74 Antibody[LN2]

Catalog Number: E-AB-F1072B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. LN₂

Biotin Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793B] Isotype Control

Conjugation

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow **FCM**

> cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µL volume or 100 µL of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD74;DHLAG;HLA class II histocompatibility antigen gamma chain;la antigen-

associated invariant chain;li;p33

Uniprot ID P04233 Gene ID 972

Background CD74 is a type II transmembrane glycoprotein also known as MHC class II associated

> invariant chain, invariant chain, li, MHC class II chaperone, and MIF receptor. CD74 exists in four isoforms with molecular masses of 33, 35, 41, and 43 kD, depending on genetic splicing. CD74 is primarily expressed on antigen presenting cells, including B cells, monocytes/macrophages, dendritic cells, and Langerhans cells. It is also expressed by activated T cells and activated endothelial and epithelial cells as well as carcinomas of lung, renal, gastric and thymic origin. The primary function of CD74 is intracellular sorting of MHC class II molecules and regulation of exogenous peptide loading onto MHC class II. It is also involved in the modulation of B cell differentiation and positive selection of CD4+ T cells. It has been reported that CD74 binds MIF (macrophage migration inhibitory factor) and signals through CD44 to regulate innate and adaptive immunity. It is also reported that H. pylori infection occurs through urease B binding of CD74 on gastric epithelial cells, inducing gastric epithelial cell apoptosis,

NF-κB activation, and IL-8 production.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com