

Recombinant Rat LTBR/TNFRSF3 Protein (His Tag)

Catalog Number: PKSR030340

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Rat
Source	HEK293 Cells-derived Rat LTBR/TNFRSF3 protein Met 1-Ala 218, with an C-terminal His
Calculated MW	23 kDa
Observed MW	35 kDa
Accession	NP_001008316.1
Bio-activity	Not validated for activity

Properties

Purity	> 96 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 96 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

LTBR (lymphotoxin beta receptor (TNFR superfamily, member 3)) is a member of the tumor necrosis factor (TNF) family of receptors. Tumor necrosis factor receptor is a trimeric cytokine receptor that binds tumor necrosis factors. The receptor cooperates with an adaptor protein (such as TRADD, TRAF, RIP), which is important in determining the outcome of the response. LTBR is expressed on the surface of most cell types, including cells of epithelial and myeloid lineages, but not on T and B lymphocytes. LTBR specifically binds the lymphotoxin membrane form (a complex of lymphotoxin-alpha and lymphotoxin-beta). LTBR and its ligand play a role in the development and organization of lymphoid tissue and transformed cells. Activation of this protein can trigger apoptosis. Not only does the LTBR help trigger apoptosis, it can lead to the release of the cytokine interleukin 8. Overexpression of LTBR in HEK293 cells increases IL-8 promoter activity and leads to IL-8 release. It is also essential for development and organization of the secondary lymphoid organs and chemokine release.