Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Mouse CD326 Antibody[G8.8]

Catalog Number: GF11810P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Immunogen Recombinant Mouse CD326 protein

Host Rat

Isotype Rat IgG2a,κ

Clone G8.8

Purification >98%, Protein A/G purified

Conjugation Unconjugated


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



4T1 were stained with 0.2 μg Purified Anti-Mouse CD326 Antibody[G8.8] (Right) and 0.2 μg Rat IgG2a, κ Isotype

Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid

freeze / thaw cycles.

Shipping Ice bag

Background

Epithelial Cell Adhesion Molecule (EpCAM), also known as GA733-2 antigen, is a type I transmembrane glycoprotein composed of an extracellular domain with two EGF-Like repeats and a cystenin-rich region, a transmembrane domain and a cytoplasmic domain. It modulates cell adhesion and proliferation. Its overexpression has been detected in many epithelial tumours and has been associated with high stage, high grade and a worse survival in some tumour types. EpCAM has been shown to function as a calcium-independent homophilic cell adhesion molecule that does not exhibit any obvious relationship to the four known cell adhesion molecule superfamilies. However, recent insights have revealed that EpCAM participates in not only cell adhesion, but also in proliferation, migration and differentiation of cells. In addition, recent study revealed that EpCAM is the Wnt-beta-catenin signaling target gene and may be used to facilitate prognosis. It has oncogenic potential and is activated by release of its intracellular domain, which can signal into the cell nucleus by engagement of elements of the wnt pathway.