

PDIA4 Polyclonal Antibody

catalog number: E-AB-18854

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human PDIA4
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:40-1:200

Data



Western blot analysis of RAW264.7 and NIH/3T3 cell lysates using PDIA4 Polyclonal Antibody at dilution of 1:250

Observed-MW:Refer to figures

Calculated-MW:73 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PDIA4 Polyclonal Antibody at dilution of

1:40(×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using PDIA4 Polyclonal Antibody at dilution of

1:40(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

For Research Use Only

PDIA4(Protein disulfide-isomerase A4) is also named as ERP70,ERP72 and belongs to the protein disulfide isomerase family. It catalyzes the rearrangement of -S-S- bonds in proteins. ERp72 is a soluble protein localized in the ER lumen and contains the COOH-terminal retention signal,KEEL. There are 6 cysteine residues in the amino acid sequences of mouse and human ERp72. All of the cysteine residues occur in the internal thioredoxin motif,CGHC. The full length protein has a signal peptide with 20 amino acid.

For Research Use Only

Toll-free: 1-888-852-8623
Web:www.elabscience.com

Tel: 1-832-243-6086
Email:techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7