# **Elabscience**®

## **BYSL Polyclonal Antibody**


### catalog number: E-AB-65860


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human BYSL (NP_004044.3).
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
	Decomposed of Dilastica


Applications	Recommended Dilution
IHC	1:50-1:100

#### Data





Immunohistochemistry of paraffin-embedded Rat ovary using Immunohistochemistry of paraffin-embedded Human colon BYSL Polyclonal Antibody at dilution of 1:100 (40x lens). carcinoma using BYSL Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human breast cancer using BYSL Polyclonal Antibody at dilution of 1:100

(40x lens).	
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.

#### Background

### For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

# **Elabscience**®

Bystin is expressed as a 2-kb major transcript and a 3.6-kb minor transcript in SNG-M cells and in human trophoblastic teratocarcinoma HT-H cells. Protein binding assays determined that bystin binds directly to trophinin and tastin, and that binding is enhanced when cytokeratins 8 and 18 are present. Immunocytochemistry of HT-H cells showed that bystin colocalizes with trophinin, tastin, and the cytokeratins, suggesting that these molecules form a complex in trophectoderm cells at the time of implantation. Using immunohistochemistry it was determined that trophinin and bystin are found in the placenta from the sixth week of pregnancy. Both proteins were localized in the cytoplasm of the syncytiotrophoblast in the chorionic villi and in endometrial decidual cells at the uteroplacental interface. After week 10, the levels of trophinin, tastin, and bystin decreased and then disappeared from placental villi.

For Research Use Only