

Recombinant Human DC-SIGN/CD209 (N-Fc)

Catalog Number:PKSH033989



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD209 molecule;CD209;CDSIGNHIV gpl20-binding protein;CLEC4L;DCSIGN;DC-SIGN;DC-SIGN1;CD209 Antigen
Species	Human
Expression Host	HEK293 Cells
Sequence	Gln59-Ala404
Accession	Q9NNX6
Calculated Molecular Weight	65.3 kDa
Observed molecular weight	75 kDa
Tag	N-Fc

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD209 is also known as CLEC4L, DC-SIGN and CD209 antigen, is a type II transmembrane protein on DCs with a C-type lectin extracellular domain, is capable of binding ICAM-3 on resting T cells in the secondary lymphoid organs, providing the initial contact between these cells during the establishment of cell-mediated immunity. The DC-SIGN/CD209 lectin domain binds mannose oligosaccharides on pathogens including HIV as well as self glycoproteins including ICAMs (2, 4). DC-SIGN/CD209 binds to butyrophilin 2A1 and this interaction can be blocked by HIV gp120. DC-SIGN/CD209 is expressed on dendritic cells (DC) and inflammatory macrophages and contributes to antigen presentation. It is not only a pattern recognition receptor but implicated in immunoregulation of DCs. It has important role

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human DC-SIGN/CD209 (N-Fc)

Catalog Number:PKSH033989



in mediating DC adhesion, migration, inflammation, activating primary T cell, triggering immune response and participating in immune escape of pathogens and tumors.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com