# Purified Anti-Human CD203c Antibody[NP4D6]

catalog number: E-AB-F12970P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Human

Immunogen Recombinant Human CD203c protein

**Host** Mouse

**Isotype** Mouse  $IgGl, \kappa$ 

Clone NP4D6

**Purification** >98%, Protein A/G purified


**Conjugation** Unconjugated

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

# **Applications** Recommended Dilution

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

#### Data



Human peripheral blood lymphocytes cell were stained with 0.2 μg Purified Anti-Human CD203c Antibody[NP4D6] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD123 PEconjugated Monoclonal Antibody.

# **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** Order now, ship in 3 days

## Background

The protein encoded by this gene belongs to a series of ectoenzymes that are involved in hydrolysis of extracellular nucleotides. These ectoenzymes possess ATPase and ATP pyrophosphatase activities and are type II transmembrane proteins. Expression of the related rat mRNA has been found in a subset of immature glial cells and in the alimentary tract. The corresponding rat protein has been detected in the pancreas, small intestine, colon, and liver. The human mRNA is expressed in glioma cells, prostate, and uterus. Expression of the human protein has been detected in uterus, basophils, and mast cells. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene.

#### For Research Use Only