

Recombinant Cynomolgus Fc gamma RIIIA/FCGR3A/CD16a (C-6His)

Catalog Number: PKSQ050099

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Cynomolgus macaques
Source	HEK293 Cells-derived Cynomolgus macaques FCGR3A/CD16a protein Glu21-Gly206, with an C-terminal His
Mol_Mass	22 kDa
Accession	A0A140HDP8
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Receptors for the Fc region of immunoglobulin G (FcγR) are divided into three classes and FcγRIII is a multifunctional, low/intermediate affinity receptor. In humans, FcγRIII is expressed as two distinct forms (FcγRIIIA and FcγRIIIB) that are encoded by two different but highly homologous genes in a cell type-specific manner. FcγRIIIB is a low-affinity, GPI-linked receptor expressed by neutrophils and eosinophils, whereas FcγRIIIA is an intermediate affinity polypeptide-anchored transmembrane glycoprotein expressed by a subset of T lymphocytes, natural killer (NK) cells, monocytes, and macrophages. The FcγRIIIA receptor is involved in phagocytosis, secretion of enzymes, inflammatory mediators, antibody-dependent cellular cytotoxicity (ADCC), mast cell degranulation, and clearance of immune complexes. FcγRIIIA has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain and delivers an activation signal in the immune responses. Aberrant expression or mutations in this gene is implicated in susceptibility to recurrent viral infections, systemic lupus erythematosus, and alloimmune neonatal neutropenia. In humans, it is a 50-70 kD type I transmembrane activating receptor. The FcγRIIIA cDNA encodes 254 amino acid including a 16aa signal sequence, 191 amino acid ECD with two C2-type Ig-like domains, five potential N-glycosylation sites, a 22 amino acid transmembrane sequence and a 25 amino acid cytoplasmic domain.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017