# AF/LE Purified Anti-Human CD18 Antibody[TS1/18.1.2.11]

catalog number: E-AB-F10570



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Human

Immunogen Recombinant Human CD18 protein

**Host** Mouse

 $\begin{tabular}{lll} \textbf{Isotype} & Mouse IgGl, \kappa \\ \textbf{Clone} & TS1/18.1.2.11 \end{tabular}$ 

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

.

Applications	Recommended Dilution

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

#### Data



Human peripheral blood lymphocytes were stained with 0.2

μg AF/LE Purified Anti-Human CD18

Antibody[TS1/18.1.2.11] (Right) and 0.2  $\mu g$  Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by FITC-conjugated Goat

Anti-Mouse IgG Secondary Antibody.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

**Shipping** Order now, ship in 3 days

### Background

CD18 is a 90-95 kD type I transmembrane protein also known as integrin  $\beta2$  subunit, LFA-1  $\beta$  subunit, and  $\beta2$  integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated  $\alpha$  chains function in adhesion and signaling in hematopoietic cells.

#### For Research Use Only