

## DMRT3 Polyclonal Antibody

catalog number: E-AB-11162

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	Recombinant protein of human DMRT3
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications

### Recommended Dilution

<b>WB</b>	1:500-1:2000
-----------	--------------

### Data



Western Blot analysis of Mouse stomach and brain tissue  
using DMRT3 Polyclonal Antibody at dilution of 1:550

**Calculated-MW:51 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

The DMRT (doublesex and mab-3 related transcription factor) genes encode a large family of transcription factors that are related to the Drosophila doublesex proteins. Expressed primarily in the gonads, DMRT proteins contain cysteine-rich DNA-binding motifs and are thought to play an important role in sexual development. DMRT3 (doublesex and mab-3 related transcription factor 3), also known as DMRTA3, is a 472 amino acid protein that contains one DM DNA-binding domain and belongs to the DMRT family. Localized to the nucleus, DMRT3 is expressed specifically in testis and is thought to regulate transcriptional events during early sexual development. The gene encoding DMRT3 maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

### For Research Use Only