A Reliable Research Partner in Life Science and Medicine

# Recombinant Human IL-21R Protein (His Tag)

Catalog Number: PKSH033365

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Species Human

Source HEK293 Cells-derived Human IL-21R protein Cys20-Pro236, with an C-terminal His

Calculated MW 26.0 kDa
Observed MW 40-55 kDa
Accession Q9HBE5

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

 Shipping
 This product is provided as lyophilized powder which is shipped with ice packs.

 Formulation
 Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

 Normally 5%
 8% trebalose mannitol and 0.01% Tween 80 are added as protectant


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

# Background

Interleukin-21 receptor is also known as IL-21 receptor; IL-21R; CD360. In humans; it is encoded by the IL21R gene. It belongs to the type I cytokine receptor family. Type 4 subfamily contains 2 fibronectin type-III domains. Interleukin-21 receptor is selectively expressed in lymphoid tissues and highly expressed in thymus and spleen. IL-21 is produced by CD4+ T cells in response to antigenic stimulation. Its action enhances antigen-specific responses of immune cells. The biological effects of IL-21 include induction of differentiation of T-cells-stimulated B-cells into plasma cells and memory B-cells. It also promotes the anti-tumor activity of CD8+ T-cells and NK cells.

#### For Research Use Only