# Recombinant Human PD-L2/CD273 Protein (His Tag)

Catalog Number: PDMH100020



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		4.0
LIPES	crir	itio n
DCS	CIII.	otion

Species Human

Source HEK293 Cells-derived Human PD-L2/CD273 protein Leu20-Pro219, with an C-terminal

His

 Mol\_Mass
 23.6 kDa

 Accession
 Q9BQ51

**Bio-activity** Not validated for activity

## **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU/mg of the protein as determined by the LAL method

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Programmed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG family. PDCD1LG2 contains one Ig-like C 2-type domain and one Ig-like V-type domain. PDCD1LG2 is highly expressed in the heart, placenta, pancreas, lung and liver, it is weakly expressed in the spleen, lymph nodes, and thymus. PDCD1LG2 is involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.

## For Research Use Only