Laminin alpha5 Polyclonal Antibody

catalog number: E-AB-31903



Note: Centrifuge before opening to ensure complete recovery of vial contents.

				ption		
	06		TO T	m	11	\mathbf{n}
v					ш	w

Reactivity Human; Mouse

Immunogen Synthesized peptide derived from the Internal region of human Laminin α-5

Host Rabbit Isotype IgG

PurificationAffinity purificationConjugationUnconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:100-1:300

Data



Western Blot analysis of RAW264.7, HepG2, A549 cells

with Laminin alpha5 Polyclonal Antibody

Observed-MV:400 kDa Calculated-MV:400 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Components of the extracellular matrix exert myriad effects on tissues throughout the body. In particular, the laminins, a family of heterotrimeric extracellular glycoproteins, affect tissue development and integrity in such diverse organs as the kidney, lung, skin, and nervous system. It is thought that laminins mediate the attachment, migration, and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components. Laminins function as heterotrimeric complexes of alpha, beta, and gamma chains, with each chain type representing a different subfamily of proteins. The protein encoded by this gene belongs to the alpha subfamily of laminin chains and is a major component of basement membranes. Two transcript variants encoding different isoforms have been found for this gene, but the full-length nature of one of them has not been determined.

For Research Use Only