

Purified Anti-Mouse TCR β Antibody[H57-597]

catalog number: E-AB-F1123A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse TCR- β protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone	H57-597
Purification	>98%, Protein A/Gpurified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 μ g/mL(1×10^5 - 5×10^5 cells)
------------	--

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μ g Purified Anti-Mouse TCR β Antibody[H57-597] (Right) and 0.2 μ g Armenian Hamster IgG, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR- β is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR- α). It is expressed on α/β TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.

For Research Use Only