Elabscience®

Purified Anti-Human CD52 Antibody[Campath-1G]

catalog number: AN004360P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human CD52 protein
Host	Rabbit
Isotype	Rabbit IgG, κ
Clone	Campath-1G
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze
	to completely remove the stabilizer prior to labeling.
Applications	Recommended Dilution
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Human peripheral blood μg Purified Anti-Hum (Right) and 0.2 μg Mo followed by Alexa Fluon IgG Se	1^{2} 10 ³ 10 ⁴ 10 ⁵ 10 ⁷ CD52 Aces Flor® 64741 A lymphocytes were stained with 0.2 an CD52 Antibody[Campath-1G] use IgG1, κ Isotype Control (Left), (\mathbb{R} 647-conjugated Goat Anti-Mouse econdary Antibody.
Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	Ice bag
Background	
CD52 also known as C	ambridge pathology antigen 1 (CAMPATH-1), is a 25-29 kD glycoprotein containing a large N-

CD52, also known as Cambridge pathology antigen 1 (CAMPATH-1), is a 25-29 kD glycoprotein containing a large Nlinked carbohydrate moiety. The actual molecule of CD52 is only 8-9 kD. It is expressed in the male reproductive tract and on virtually all lymphocytes (T and B cells), as well as macrophages/monocytes, eosinophils, and red cells. CD52 is thought to play a role in carrying and orienting carbohydrates. CD52 is a potent target for complement-mediated lysis and antibody-mediated cellular cytotoxicity and has been used as a depletion target for chronic lymphocytic leukemia (CLL)/lymphoma and immunosuppression.