

# Recombinant Human IL-1R8/IL1RAPL1 Protein (His Tag)

Catalog Number: PKSH033634



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Human
<b>Mol_Mass</b>	40 kDa
<b>Accession</b>	Q9NZN1
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Interleukin-1 receptor accessory protein-like 1; also known as IL1RAPL1; can be detected at low levels in heart; skeletal muscle; ovary; skin; amygdala; caudate nucleus; corpus callosum; hippocampus; substantia nigra and thalamus. IL1RAPL1 functions as a homodimer; it interacts with NCS1; PTPRD. This interaction is PTPRD-splicing-dependent and induces pre- and post-synaptic differentiation of neurons and is required for IL1RAPL1-mediated synapse formation. During dendritic spine formation; it can bidirectionally induce pre- and post-synaptic differentiation of neurons by trans-synaptically binding to PTPRD.

## For Research Use Only