# AF/LE Purified Anti-Mouse CD54 Antibody[YN1/1.7.4]

catalog number: E-AB-F10180



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-				
	AC	cri	nti	nn
$\mathbf{L}$		CII		

**Reactivity** Mouse

Immunogen Recombinant Mouse CD54 protein

Host Rat

IsotypeRat IgG2b,  $\kappa$ CloneYN1/1.7.4

**Purification** >98%, Protein A/G purified


Conjugation None (AF/LE)

**buffer** Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

Applications	Recommended Dilution		

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

### Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg AF/LE Purified Anti-Mouse CD54 Antibody[YN1/1.7.4] (Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

## Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

**Shipping** Ice bag

# Background

CD54 is a 90 kD immunoglobulin superfamily member also known as ICAM-1 and Ly-47. It is expressed on activated endothelial cells, high endothelial venules (HEV), T and B cells, monocytes/ macrophages, granulocytes, and dendritic cells. CD54 is an important intracellular adhesion molecule that participates in T cell-T cell, T cell-B cell, and T cell-target cell interactions via binding of LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). CD54 has also been shown to be involved in lymphocyte trafficking, making it an important molecule in many immune reactions and inflammation. CD54 is also a receptor for rhinovirus. The YN1/1.7.4 antibody has been reported to block binding of mouse CD54 to LFA-1 and Mac-1, inhibit cell-cell adhesion, and function in antigen presentation to T cells and leukocyte migration to inflammatory tissues.

### For Research Use Only