

Recombinant Human TMED1 Protein (His Tag)

Catalog Number: PKSH030660

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human TMED1 protein Met 1-Asn194, with an C-terminal His
Calculated MW	20.6 kDa
Observed MW	28 kDa
Accession	Q13445
Bio-activity	Measured by its binding ability in a functional ELISA. 2. Immobilized human TMED1-His at 10µg/mL (100µL/well) can bind human IL1R4-Fc, the EC ₅₀ of human IL1R4-Fc is 8-50ng/mL.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

TMED1 belongs to the EMP24/GP25L family. It contains 1 GOLD domain and is widely expressed. TMED1 binds to its receptor IL1RL1 and results in the activation of DNA binding by nuclear factor NF-kappa-B or transcription from the IL8 promoter and most likely requires other proteins to elicit these activities. Dendritic cells from Peyer's patches (but not from spleen) express TMED1 in response to treatment with LPS. TMED1 may play a role in vesicular protein trafficking; mainly in the early secretory pathway. It may act as a cargo receptor at the luminal side for incorporation of secretory cargo molecules into transport vesicles and may be involved in vesicle coat formation at the cytoplasmic side.

For Research Use Only