A Reliable Research Partner in Life Science and Medicine

Recombinant Human Rad51 protein (His Tag)

Catalog Number: PDEH100994

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human Rad51 protein Ala2-Asp339, with an N-terminal His

 Mol_Mass
 37.1 kDa

 Accession
 Q06609

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Rad51 proteins, 2µg/lane of Recombinant Human Rad51 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 37 KD.

Background

Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination. Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template. Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:26681308). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3.

For Research Use Only

Fax: 1-832-243-6017