

## Recombinant Human MESDC2/MESD Protein (His Tag)

**Catalog Number:** PKSH031340

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human MESDC2/MESD protein Ala 34-Lys 230, with an C-terminal His
<b>Calculated MW</b>	23.6 kDa
<b>Observed MW</b>	27 kDa
<b>Accession</b>	NP_055969.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

LDLR chaperone MESD, also known as Mesoderm development protein, Mesoderm development candidate 2, Renal carcinoma antigen NY-REN-61 and MESDC2, is a member of the MESD family. MESDC2 is a chaperone specifically assisting the folding of beta-propeller/EGF modules within the family of low-density lipoprotein receptors (LDLRs). The LDLR maturation activity resides in the N- and C-terminal unstructured regions. MESDC2 acts as a modulator of the Wnt pathway, since some LDLRs are coreceptors for the canonical Wnt pathway. MESDC2 is essential for specification of embryonic polarity and mesoderm induction.

### For Research Use Only