Elabscience®

PCK1 Polyclonal Antibody

catalog number: D-AB-10224L


Note: Centrifuge before opening to ensure complete recovery of vial contents.


1:100-1:200

| Description | |
|--------------|--|
| Reactivity | Human;Rat |
| Immunogen | Recombinant Rat PCK1 protein expressed by E.coli |
| Host | Rabbit |
| Is otype | IgG |
| Purification | Antigen Affinity Purification |
| Conjugation | Unconjugated |
| Buffer | PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4 |
| Applications | Recommended Dilution |

IHC


Data





Immunohistochemistry of paraffin-embedded Rat kidney using Pck1 Polyclonal Antibody at dilution of 1:100

Immunohistochemistry of paraffin-embedded Human kidney using Pck1 Polyclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffin-embedded Human liver using PCK1 Polyclonal Antibody at dilution of 1:200

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |
| | |

Background

This gene is a main control point for the regulation of gluconeogenesis. The cytosolic enzyme encoded by this gene, along with GTP,catalyzes the formation of phosphoenolpyruvate from oxaloacetate,with the release of carbon dioxide and GDP. The expression of this gene can be regulated by insulin,glucocorticoids,glucagon,cAMP,and diet. Defects in this gene are a cause of cytosolic phosphoenolpyruvate carboxykinase deficiency. A mitochondrial isozyme of the encoded protein also has been characterized.