Recombinant Human ITGA5 Protein (His Tag)

Catalog Number: PKSH032600

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Human	
Source	HEK293 Cells-derived Human ITGA 5 protein Phe42-Tyr995, with an C-terminal His	
Calculated MW	105.1 kDa	
Observed MW	130 kDa	
Accession	P08648	
Bio-activity	Not validated for activity	
Properties		
Purity	> 90 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

kDa	MK	° R
120 90	=	-
60		
40	-	colence
30	-	Elalos
20	-	
14	-	

> 90 % as determined by reducing SDS-PAGE.

Background

Integrin α -5 belongs to the Integrin α chain family and contains 7 FG-GAP repeats. Integrin α -5 joins with Integrin- β 1 to form a fibronectin and laminin receptor which recognizes the sequence R-G-D in its ligands. In case of HIV-1 infection; the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions. It is expressed on fibroblasts; endothelial cells; peripheral T cells and platelets. Integrin α -5 undergoes post-translational cleavage in the extracellular domain to yield disulfide-linked light and heavy chains. In addition to adhesion; ITGA5 participates in cell-surface mediated signalling.