## **Elabscience**®

## Recombinant Human CD32b/FCGR2B Protein (HEK293 Cells, His Tag)

## Catalog Number: PKSH031726

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Human		
Source	HEK293 Cells-derived Human CD32b/FCGR2B protein Ala 46-Pro 217, with an C-		
	terminal His		
Calculated MW	20.8 kDa		
Observed MW	25-30 kDa		
Accession	NP_001002274.1		
<b>Bio-activity</b>	Immobilized recombinant human CD32b at 10 $\mu$ g/ml (100 $\mu$ l/well) can bind human		
	IgG2 with a linear range of 0.16-6. 4 $\mu$ g/ml.		
Properties			
Purity	>95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from sterile PBS, pH 7.4		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		
Data			
KDa	a MK R		
112			

кDa	IVIN	ent
116	- Harris	12.
66.2	-	clabscience
	1000	bscler
45.0	1000	Elan
35.0	-	
	1000	100
25.0	-	- apscient
	1000	F.
184	-	
	CIE	
14.4	-	
	116	116 66.2 45.0 35.0 25.0 18.4

> 95 % as determined by reducing SDS-PAGE.

## Background

Processing of the N-terminal initiator methionine or formylated methionine is an essential cellular process conserved from prokaryotes to eukaryotes. The proteolytic removal of N-terminal methionine from nascent peptides is catalyzed by a family of enzymes known as methionine aminopeptidases (MetAPs) and is essential for cell growth. METAP1 and METAP2 have different substrate specificity due to the differences in both size and shape of the active sites. As a member of the M24 family of metalloproteases, METAP1 plays an important role in G(2)/M phase regulation of the cell cycle and may serve as a promising target for the discovery and development of new anticancer agents.

For Research Use Only