A Reliable Research Partner in Life Science and Medicine

ATF1 Polyclonal Antibody

catalog number: E-AB-70146

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen KLH conjugated Synthetic peptide corresponding to Mouse ATF1

Host Rabbit Isotype IgG

Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein

protectant and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:1000

Data



Western Blot analysis of various samples using ATF1

Polyclonal Antibody at dilution of 1:1000.

Observed-MW:29 kDa Calculated-MW:29 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes an activating transcription factor, which belongs to the ATF subfamily and bZIP (basic-region leucine zipper) family. It influences cellular physiologic processes by regulating the expression of downstream target genes, which are related to growth, survival, and other cellular activities. This protein is phosphorylated at serine 63 in its kinase-inducible domain by serine/threonine kinases, cAMP-dependent protein kinase A, calmodulin-dependent protein kinase I/II, mitogen- and stress-activated protein kinase and cyclin-dependent kinase 3 (cdk-3). Its phosphorylation enhances its transactivation and transcriptional activities, and enhances cell transformation. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in angiomatoid fibrous histiocytoma and clear cell sarcoma. This gene has a pseudogene on chromosome 6.

For Research Use Only

Rev. V1.7