

## Recombinant Histone H2A (Acetyl Lys5) Monoclonal Antibody

catalog number: **AN301414L**


**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Human;Rat;Mouse
<b>Immunogen</b>	Acetylated human histone H2A (Lys5) peptide
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG, $\kappa$
<b>Clone</b>	A109
<b>Purification</b>	Protein A purified
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications

<b>WB</b>	1:1000-1:2000
<b>ChIP</b>	6 $\mu$ g/ $5 \times 10^6$ cells



Chromatin immunoprecipitation analysis of HeLa+TSA (+) (400nM, 16h) immunoprecipitated DNA by real-time PCR using primers specific for the human GAPDH-CDS, LDHA, FOXO3a-D and TUBBP10. The data are presented as enrichment of each sample relative to the total amount of input chromatin at each amplicon.

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

Histone post-translational modifications (PTMs), known as the “histone code”, are key mechanisms of epigenetics that modulate chromatin structures. The PTMs on histone including acetylation, methylation, phosphorylation, and novel acylations directly affect the accessibility of chromatin to transcription factors and other epigenetic regulators, altering genome stability and gene transcription. Histone acetylation, tightly controlled by the opposing action of histone acetyltransferases (HATs) and histone deacetylases (HDACs), occurs primarily at lysine residues on the N-terminal tails of histones H2A (Lys5, 9, and 15), H2B (Lys5, 12, 15, 16, and 20), H3 (Lys4, 9, 14, 18, 23, 27, and 36), and H4 (Lys5, 8, 12, 16, and 20), and plays vital roles in the regulation of gene expression, DNA damage repair, chromatin dynamics, etc.

### For Research Use Only