

Recombinant Human TIMP2/TIMP-2 Protein (His Tag)

Catalog Number: PKSH033118

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human TIMP2;TIMP-2 protein Cys27-Pro220, with an C-terminal His
Calculated MW	22.8 kDa
Observed MW	26 kDa
Accession	P16035
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). There are four members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4. Tissue Inhibitor of Metalloproteinases 2 (TIMP-2) is a non N-glycosylated protein with a molecular mass of 22 kDa. It produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes and irreversibly inactivates them by binding to their catalytic zinc cofactor. TIMP-2 also has erythroid-potentiating and cell growth promoting activities.

For Research Use Only