Recombinant Human Interleukin-2/IL-2 Protein (C145S)

Catalog Number: PKSH033623



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 10	AC	cri	m		nn
\mathbf{L}			ш	UΙU	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 15.5 kDa

 Accession
 P60568

Bio-activity Measured in a cell proliferation assay using CTLL- 2 mouse cytotoxic T cells. The

ED50 for this effect is 90-270 pg/ml

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.
 Formulation Lyophilized from a 0.2 μm filtered solution of 10mM Acetata-Na, 5% Trehaiose, pH

4.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.

For Research Use Only