

TRIP4 Polyclonal Antibody

catalog number: E-AB-14750

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Immunogen	Recombinant protein of human TRIP4
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution


WB	1:500-1:2000
IHC	1:25-1:100

Data




Antibody at dilution of 1:400

Calculated-MW: 66 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer using TRIP4 Polyclonal Antibody at dilution of 1:50



Immunohistochemistry of paraffin-embedded Human liver cancer using TRIP4 Polyclonal Antibody at dilution of 1:50

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Activating signal co-integrator-1 (ASC-1), originally identified as TRIP4, is a transcriptional co-activator of nuclear receptors that associates with specific components of the RNA polymerase II complex and binds the basal transcription factors TBP and TFIIA. ASC-1 functions with the transcription integrators SRC-1 and CBP/p300 through its zinc finger motif and is dependent on their ligand-dependent transactivation domain, AF2. Endogenous ASC-1 in HeLa cells is predominantly a nuclear protein. Under conditions of serum starvation, ASC-1 localizes to the

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8