Recombinant Human KDR Protein(Fc Tag)

Catalog Number: PDMH100192



Note: Centrifuge before opening to ensure complete recovery of vial contents.

	escrip				7.0		
	00		TO T	m	TT.		т
v		ж			ш	w	٠

Species Human

Source Mammalian-derived Human KDR protein Ala20-Glu764, with an C-terminal Fc

 Mol_Mass
 106.8 kDa

 Accession
 P35968

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human KDR proteins, 2 µg/lane of Recombinant Human KDR proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 106.8

KD

Background

Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

For Research Use Only