

# Recombinant Human ENPP7/NPP-7 Protein (His Tag)

Catalog Number:PKSH031375



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	ALK-SMase;E-NPP7;NPP-7;NPP7
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Ser 439
<b>Accession</b>	NP_848638.2
<b>Calculated Molecular Weight</b>	49.0 kDa
<b>Observed molecular weight</b>	55-60 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Ectonucleotide pyrophosphatase / phosphodiesterase family member 7, also known as Alkaline sphingomyelin phosphodiesterase, Intestinal alkaline sphingomyelinase, Alk-Smase, ENPP7 and NPP-7, is a single-pass type I membrane protein which belongs to thenucleotide pyrophosphatase / phosphodiesterase family. ENPP7 / NPP-7 is expressed in the intestines and human bile. ENPP7 / NPP-7 is localized at the surface of the microvillar membrane in small intestine enterocytes, as well as in endosome-like structures and in Golgi complex. The main function of ENPP7 / NPP-7 is to convert the dietary sphingomyelin into ceramide, the sphingolipid messengers via hydrolyzation. ENPP7 / NPP-7 is also reported to exert a phospholipase C activity toward palmitoyl lyso-phosphocholine. The activity of this enzyme is inhibited in a dose dependent manner by ATP, imidazole, orthovanadate and zinc ion. Further, It has been shown in

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human ENPP7/NPP-7 Protein (His Tag)

Catalog Number:PKSH031375



studies that decreased levels of ENPP7 / NPP-7 may be associated with human colon cancer.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)