

## Recombinant Human Vitamin D-Binding Protein/GC protein (His Tag)

Catalog Number: PDMH100111

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Vitamin D-Binding;GC protein Met1-Leu474, with an C-terminal His
<b>Calculated MW</b>	52.0 kDa
<b>Observed MW</b>	57 kDa
<b>Accession</b>	P02774
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Vitamin D-Binding Protein (DBP) is a member of the ALB/AFP/VDB family. DBP is a secreted protein and contains three albumin domains. The primary structure contains 28 cysteine residues forming multiple disulfide bonds. DBP acts as a multifunctional protein found in plasma, ascitic fluid, cerebrospinal fluid, and urine and on the surface of many cell types. DBP binds to vitamin D and its plasma metabolites and transports them to target tissues. DBP associates with membrane-bound immunoglobulin on the surface of B-lymphocytes and with IgG Fc receptor on the membranes of T-lymphocytes.

### For Research Use Only