

Rat IL-6 Antibody Pair Set

Catalog No.	E-KAB-0106	Applications	ELISA
Synonyms	IL6, BSF2, HGF, HSF, IFNB2, CDF		

Kit components & Storage

Title	Specifications	Storage
Rat IL-6 Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Rat IL-6 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0106)	
		Rat IL-6 Capture Antibody	Rat IL-6 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Rat IL-6 protein	Recombinant Rat IL-6 protein
	Swissprot	P20607	
Product details	Reactivity	Rat	Rat
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Rat IL-6 in ELISAs.	

Applications

Rat IL-6 Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4µg/mL	Rat IL-6 Capture Antibody	
ELISA Detection	1:1000-1:10000	Rat IL-6 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Cytokine with a wide variety of biological functions in immunity, tissue regeneration, and metabolism. Binds to IL6R, then the complex associates to the signaling subunit IL6ST/gp130 to trigger the intracellular IL6-signaling pathway (Probable). The interaction with the membrane-bound IL6R and IL6ST stimulates 'classic signaling', whereas the binding of IL6 and soluble IL6R to IL6ST stimulates 'trans-signaling'.

Alternatively, 'cluster signaling' occurs when membrane-bound IL6:IL6R complexes on transmitter cells activate IL6ST receptors on neighboring receiver cells (Probable). IL6 is a potent inducer of the acute phase response. Rapid production of IL6 contributes to host defense during infection and tissue injury, but excessive IL6 synthesis is involved in disease pathology. In the innate immune response, is synthesized by myeloid cells, such as macrophages and dendritic cells, upon recognition of pathogens through toll-like receptors (TLRs) at the site of infection or tissue injury (Probable). In the adaptive immune response, is required for the differentiation of B cells into immunoglobulin-secreting cells. Plays a major role in the differentiation of CD4+ T cell subsets. Essential factor for the development of T follicular helper (Tfh) cells that are required for the induction of germinal-center formation. Required to drive naive CD4+ T cells to the Th17 lineage. Also required for proliferation of myeloma cells and the survival of plasmablast cells.

Acts as an essential factor in bone homeostasis and on vessels directly or indirectly by induction of VEGF, resulting in increased angiogenesis activity and vascular permeability.

Induces, through 'trans-signaling' and synergistically with IL1B and TNF, the production of VEGF.

Involved in metabolic controls, is discharged into the bloodstream after muscle contraction increasing lipolysis and improving insulin resistance.

'Trans-signaling' in central nervous system also regulates energy and glucose homeostasis.

Mediates, through GLP-1, crosstalk between insulin-sensitive tissues, intestinal L cells and pancreatic islets to adapt to changes in insulin demand.

Also acts as a myokine (Probable). Plays a protective role during liver injury, being required for maintenance of tissue regeneration.

Also has a pivotal role in iron metabolism by regulating HAMP/hepcidin expression upon inflammation or bacterial infection.

For Research Use Only

Through activation of IL6ST-YAP-NOTCH pathway, induces inflammation-induced epithelial regeneration.