# Purified Anti-Human CD235a Antibody[10F7MN]

catalog number: AN004100P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

**Reactivity** Human

Immunogen Recombinant Human CD235a protein

**Host** Mouse

IsotypeMouse IgGl,  $\kappa$ Clone10F7MN

**Purification** >98%, Protein A/G purified


**Conjugation** Unconjugated

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

## **Applications** Recommended Dilution

FCM  $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ 

#### Data



Human peripheral blood were stained with 0.2 μg Purified Anti-Human CD235a Antibody[10F7MN] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD45 FITC-conjugated Monoclonal Antibody and anti-Human CD61 PE-conjugated Monoclonal Antibody. Analyzed CD45 negative and CD61 positive populations.

#### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

#### **Background**

CD235a (Glycophorin A) is member of the glycophorin A family. It is a type I sialoglycoprotein with a molecular weight of 10 kD, present in the cell membrane as a homodimer. Glycophorin A is expressed by erythroid precursors and erythrocytes. It carries the antigen determinants for the MNS blood groups and has been proposed to be an inhibitor of hemagglutination and hemolysis. Glycophorin A binds siglec 5, the erythrocyte binding antigen (EBA-175) of P. falciparum and some viruses, including influenza virus and hepatitis A virus.

# For Research Use Only