

## Recombinant Human SUV420H2 Protein (His & GST Tag)

**Catalog Number:** PKSH030951

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human SUV420H2 protein Gly 2-Leu 280, with an N-terminal His & GST
<b>Calculated MW</b>	60.0 kDa
<b>Observed MW</b>	60 kDa
<b>Accession</b>	NP_116090.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 80 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 50mM Tris, 0.5M NaCl, 30% Glycerol, 0.05% Tween, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 80 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Histone-lysine N-methyltransferase SUV420H2, also known as Suppressor of variegation 4-20 homolog 2, Su(var)4-20 homolog 2, Lysine N-methyltransferase 5C, SUV420H2 and KMT5C, is nucleus protein which belongs to the histone-lysine methyltransferase family and Suvar4-20 subfamily. SUV420H2 is a histone methyltransferase that specifically trimethylates 'Lys-20' of histone H4. H4 'Lys-20' trimethylation represents a specific tag for epigenetic transcriptional repression. SUV420H2 mainly functions in pericentric heterochromatin regions, thereby playing a central role in the establishment of constitutive heterochromatin in these regions. SUV420H1 is targeted to histone H3 via its interaction with RB1 family proteins (RB1, RBL1 and RBL2). FRAP experiments reveal that SUV420H2 is strongly associated to pericentric heterochromatin. The fraction of SUV420H2 captured and characterized by TAP/MS is a soluble fraction which may be in a stable association with HP1. SUV420H2 may be recruited to heterochromatin in association with HP1, and stably maintained at its heterochromatin sites in an HP1-independent fashion.