

A Reliable Research Partner in Life Science and Medicine

Elab Bright™ Violet 510 Anti-Mouse CD4 Antibody[GK1.5]

Catalog Number: E-AB-F1097R1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2b, κ **Clone No.** GK1.5

Isotype Control Elab Bright™ Violet 510 Rat IgG2b, κ Isotype Control[R35-38] [Product AN00821R1]

Conjugation Elab Bright™Violet 510

Conjugation Information Elab BrightTM Violet 510 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 510 nm (e.g., a 525/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with Elab

BrightTM Violet 510 Ms CD4 GK1.5 50ug (left) or Elab Bright TM Violet 510 Rat IgG2b, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4

 Uniprot ID
 P06332

 Gene ID
 12504

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosin kinase, Ick.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com